



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Hawks took possession for breeding purposes, of an apartment in a dove-cote at my farm in Tatham in West Springfield, driving out a pair of Doves that were there in possession and destroying their nest. Their first egg was laid April 17, the second after an interval of two days, and three others, each, after an interval of one day. Incubation commenced after the fourth was laid. The male was at this time killed, but the female remained devoted to her work and on the 27th of May three Hawks were hatched, and the following day, another. One of the eggs proved not to be fertile. Incubation lasted thirty-four days, a period much longer than heretofore reported. During the whole of the time of incubation and the rearing of the young, the mother Hawk did not interfere with the wild birds that had adopted the territory in the vicinity of the dove-cote for their home.

A pair of Bluebirds nested in a bird-house within thirty feet, and Robins, Phœbes, Vesper Sparrows and other kinds all remained undisturbed in the immediate neighborhood, and the pair of Doves that were first made to give way for the Hawks, were permitted to rebuild in a place adjacent to their first home. The young of the Hawks were all successfully raised and are now well and happy in confinement.—ROBERT O. MORRIS, *Springfield, Mass.*

**Xema sabinii** and **Chordeiles virginianus sennetti**—Two Additions to the Iowa Avifauna. — My collection of Iowa birds contains two immature specimens of Sabine's Gull, both of which were taken on the sandbar immediately above Burlington, Iowa. No. 50, (S. U. I. No. 15981) ♂, was shot Oct. 15, 1891; No. 51, (S. U. I. No. 15982) ♀, Oct. 12, 1894. These I believe are the first records of this species for Iowa. The specimens are deposited at the State University of Iowa at Iowa City.

The Smithsonian Institution recently received a specimen of Sennett's Night Hawk from Mr. C. F. Henning of Boone, Iowa, shot four miles southeast of that place. This variety seems so far to have escaped Iowa observers and it gives me pleasure to add it to our list.—P. BARTSCH, *Washington, D. C.*

---

## RECENT LITERATURE.

**Torrey's 'A World of Green Hills.'**<sup>1</sup>—The subtitle of Mr. Torrey's

---

<sup>1</sup> A World of Green | Hills | Observations of Nature | and Human Nature | in the Blue Ridge | By | Bradford Torrey | . . . [Motto, = 2 lines and Seal] Boston and New York | Houghton, Mifflin and Company | The Riverside Press, Cambridge | 1898 — 16mo, pp. 285. Price, \$1.25.